NOAA Technical Report NMFS 29

Prevalence, Intensity, Longevity, and Persistence of *Anisakis* sp. Larvae and *Lacistorhynchus tenuis* Metacestodes in San Francisco Striped Bass

Mike Moser, Judy A. Sakanari, Carol A. Reilly, and Jeannette Whipple

April 1985

LIBRARY
APR 19 1985
National Marine Fisheries Service
2570 Dole Street
Honolulu, Hawaii

NOAA
TEch.
Rept.
NMFS
29

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS, intensive scientific reports on studies of restricted scope, papers on applied fishery problems, technical reports of general interest intended to aid conservation and management, reports that review in considerable detail and at a high technical level certain broad areas of research, and technical papers originating in economics studies and from management investigations.

Copies of NOAA Technical Report NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.

NMFS Honolulu Library
00000000004716

NOAA Technical Report NMFS 29

Prevalence, Intensity, Longevity, and Persistence of *Anisakis* sp. Larvae and *Lacistorhynchus tenuis* Metacestodes in San Francisco Striped Bass

Mike Moser, Judy A. Sakanari, Carol A. Reilly, and Jeannette Whipple

April 1985

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
John V. Byrne, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator for Fisheries

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any propriety product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction 1
Materials and methods 1
  Collection 1
  Aging 1
  Diet analysis 1
  Statistical analysis 1
  Serial infections 1
  Longevity and persistence 2
Results 2
  Diet analysis 2
  Anisakis sp. 2
    Host reaction 2
    Prevalence and intensity 2
    Serial infection 2
    Longevity and persistence 3
  Lacistorhynchus tenuis 3
    Host reaction 3
    Prevalence and intensity 3
    Persistence 3
  Other parasites and a tumor 3
Discussion 3
Acknowledgments 4
Literature cited 4

Figures

1. Stomach content analysis of striped bass in the three major age groups in winter, spring, and summer 2
2. Cross section of posterior end of stomach with embedded larval nematodes 2
3. Prevalence and intensity of larval nematodes compared with age of host 2
4. *Anisakis* sp. larvae encapsulated in fibrous "raft" 3
5. Prevalence and intensity of metacestodes compared with age of host 3
6. Tumor, attached to mesenteries, cross section 3



Digitized by the Internet Archive
in 2025

https://archive.org/details/noaa-technical-report-nmfs_1985-04_29

Prevalence, Intensity, Longevity, and Persistence of *Anisakis* sp. Larvae and *Lacistorhynchus tenuis* Metacestodes in San Francisco Striped Bass

MIKE MOSER, JUDY A. SAKANARI, CAROL A. REILLY,[1] and JEANNETTE WHIPPLE[2]

ABSTRACT

Thirteen hundred and seventy-three striped bass, *Morone saxatilis*, were collected from the San Francisco Bay-Delta area to correlate host diet with parasitic infections and to determine the prevalence, intensity, longevity, and persistence of larval *Anisakis* sp. nematodes and the metacestode *Lacistorhynchus tenuis*. There is an increase in the prevalence and intensity of *Anisakis* sp. and in the intensity of *L. tenuis* with increase of age of the host. These increases are probably related to the diet and the persistence of the parasites. The infections of both species are overdispersed. San Francisco Bay striped bass are an incompatible host for both species of parasites. Degenerated *Anisakis* sp. will remain in the host for at least 8 months and *L. tenuis* metacestodes for 22 months. The occurrence of several other species of parasites and a tumor are also reported.

INTRODUCTION

Striped bass, *Morone saxatilis*, is a popular sportfish in the San Francisco Bay, CA, area. Anglers have often reported the presence of parasites in the mesenteries and muscle of striped bass and questioned the public health aspects of the parasitic infections. The purpose of this study was to determine the prevalence (no. hosts infected/no. hosts examined), intensity (total no. parasites/no. infected hosts), longevity (time parasites remain alive), and persistence (time degenerated parasites are detectable) of the two most commonly seen parasites: the metacestode *Lacistorhynchus tenuis* (Cestoda) and the larval nematode *Anisakis* sp. (probably *A. simplex*).

MATERIALS AND METHODS

Collection

Thirteen hundred and seventy-three striped bass were collected from the San Francisco Bay-Delta area between June 1979 and December 1983 by hook and line, gill net, and otter trawl. Fish were frozen or Formalin-fixed and autopsied for parasites and stomach content analyses at Long Marine Laboratory, University of California, Santa Cruz.

Aging

The age of the fish was determined by scale ring counts by the California Department of Fish and Game and the use of growth curves.

Diet Analysis

The stomach contents of the fish were analyzed to determine possible correlations between diet and parasite infections. Stomach contents of 831 striped bass collected between January and September 1981 from San Pablo Bay and Suisun Slough were examined for possible intermediate hosts. The Index of Relative Importance (IRI) of food items was determined by methods suggested by Cailliet (1976). The young-of-the-year (YOY) and most of the 1-yr-olds had similar diets, hence were combined for analysis. The age groups for diet analysis were: YOY and 1-yr-old (10 to 230 mm SL); subadult females and adult males (2 to 3 yr); and adults (4+ yr).

Statistical Analysis

The coefficient of dispersion (overdispersed distribution) was determined to indicate possible clumping in parasite distributions, hence the possible need for nonparametric analysis (Sokal and Rohlf 1969). Within each age class seasonality did not significantly change the prevalance or intensity of the parasitic infections. Therefore, all seasons were combined within each age class (ANOVA, regression analysis).

Serial Infections

Serial infections were performed to determine whether nematode larvae can be transferred from fish to fish. The striped bass used in both infection experiments were YOY hatchery fish provided by the California Department of Fish and Game, Elk Grove, CA. The third stage larval *Anisakis* sp. used were from bocaccio, *Sebastes paucispinis*. The fish were force-fed (Wootten and Smith 1975) with washed larvae (Tanaka and Torisu 1978). Fifty larvae were fed to each of six striped bass. As a control, four YOY yellowtail rockfish, *Sebastes flavidus*, were infected in a similar manner.

[1]Coastal Marine Studies, University of California, Santa Cruz, Santa Cruz, CA 95064.

[2]Physiological Ecology Investigations, Southwest Fisheries Center Tiburon Laboratory, National Marine Fisheries Service, NOAA, Tiburon, CA 94920.

Longevity and Persistence

Striped bass collected from the field were observed to have degenerated larval nematodes in their mesenteries and tissues. The purposes of this experiment were to determine how long the larval *Anisakis* sp. remain alive in striped bass, therefore potentially harmful to humans, and to determine the persistence of dead larvae in the mesenteries and muscle. The third stage *Anisakis* sp. larvae used were prepared as above but were from sablefish, *Anoplopoma fimbria*. To insure that all fish were infected with equal numbers of larvae, 25 larvae were inserted into the coelomic cavity through a small cut on the sides of 178 striped bass. This wound was sealed with a nontoxic biological glue, Histoacryl (Tri-Hawk, Los Angeles, CA). Previous tests showed that striped bass reacted similarly to larvae infected by this technique, to those force-fed, and to those seen in field fish. The hosts were periodically autopsied during a 33-wk period.

RESULTS

Diet Analysis

The fullness of the stomachs and the percent IRI for the dominant food item or items are given for each age class. No stomachs contained larvae or metacestodes (Fig. 1).

Anisakis sp.

Host reaction.—The larval *Anisakis* from field fish were found on the mesenteries and internal organs. Rarely were they found in the muscle tissue. Most larvae were dead or degenerating with dark brown cuticles. The larvae were often seen as solid, round masses of fibrous tissue. When several larvae were in close proximity in the mesenteries, they were often encapsulated to form a fibrous "raft." In field fish with high intensity infections, numerous degenerated larvae were found embedded in the posterior end of the stomach (Fig. 2).

Prevalance and intensity.—The prevalence and intensity of larval nematode infections in striped bass begin to increase at 4 yr and continue to increase throughout the life of the host. The mean to variance ratios (year classes 3 to 8+ are: 1.3, 14.2, 58.8, 33.7, 49.9, 39.5, respectively) show the infections to be increasingly overdispersed with time (Fig. 3).

Serial infection.—Autopsies showed that after 5 wk, two of six striped bass were infected by one and four larvae, respectively.

Figure 2.—Cross section of posterior end of stomach with embedded larval nematodes.



Figure 3.—Prevalence and intensity of larval nematodes compared with age of host. Bar = standard error.

Although the larvae from the striped bass responded weakly to tactile stimuli, the cuticle had darkened, resembling the dead larvae recovered from field-captured striped bass. In contrast, three of the four rockfish were infected with an average of four larvae. The larvae from the rockfish were more active, and the cuticle was flesh colored, similar to larvae from field-collected *Sebastes paucispinis* and other fish hosts.



Figure 1.—Stomach content analysis of striped bass in the three major age groups in winter (W), spring (Sp), and summer (Su).

Longevity and persistence.—At the end of 2 wk most larvae had been encapsulated into a fibrous "raft." In the following 31 wk, in addition to the "raft," one to five larvae were recovered in the coelom or attached to the mesenteries. These larvae usually responded weakly to tactile stimuli, but became progressively darker and weaker with time (Fig. 4).

Figure 4.—*Anisakis* sp. larvae encapsulated in fibrous "raft."

Lacistorhynchus tenuis

Host reaction.—Moser, Sakanari, Wellings, and Lindstrom (1984) described the inflammatory response of the striped bass to these "rafts." The metacestodes from field fish appeared degenerated. The outer surface of the blastocyst was brown and the metacestode was a hardened mass. Often, the metacestodes, similar to the nematode larvae, were encapsulated into fibrous "rafts."

Prevalance and intensity.—During the first year the prevalence of larval cestode infections reached 81.6% and remained approximately at this level throughout the year. However, the mean intensity of infection slowly increases with the age of the host. The mean to variance ratios (age classes YOY to 8+ are: 14.4, 23.2, 38.9, 9.2, 48.6, 36.7, 59.3, 51.2, 73.0, respectively) indicate the infections were increasingly overdispersed with time (Fig. 5).

Persistence.—Four field fish kept in the laboratory for 22 mo still retained their dead metacestodes and "rafts."

Other Parasites and a Tumor

Occasionally other helminth parasites were encountered during autopsies. However, because of the overall goal of this study and the method of preservation of field fish, no attempt was made to quantify these helminths. Parasites reported from the San Francisco Bay-Delta include: Acanthocephala—*Southwellina* (D. Woodbury[3]); Cestoda—Tetraphyllidae larvae, in mesenteries,

[3]D. Woodbury, Seasonal Aide, California Department of Fish and Game, 203 Tall Trees Drive, Vallejo, CA 94589, pers. commun. March 1983.



Figure 5.—Prevalence and intensity of metacestodes compared with age of host. Bar = standard error.

(this study); Digenea—*Stephanostomum tenue* (D. Woodbury footnote 3), *Lepocreadium californianum* (Edwards and Nahhas 1968); Monogenea—*Cleidodiscus pricei* (Hensley and Nahhas 1975); Nematoda—*Thynnascaris brachyurum* (as *Contracaecum b.*, Hensley and Nahhas 1975), *Spiroxys* sp. larva (Hensley and Nahhas 1975), and *Phocanema* sp. larvae, in mesenteries, (this study); a tumor, in coelom, (this study, Fig. 6, National Museum of Natural History-Registry of Tumors in Lower Animals (NMNH-RTLA) #3173).



Figure 6.—Tumor a) attached to mesenteries, b) cross section.

DISCUSSION

The food habits of San Francisco Bay-Delta striped bass have been reported by Turner and Kelley (1966), Thomas (1967), Orsi (1971), Moriguchi (1973), Moyle (1976), and Orsi and Knutson (1979). The present study and the above reports show that YOY consume primarily shrimp (*Neomysis* sp. and *Crangon* sp.), amphipods (*Corophium* sp.), and copepods. By the end of the first year the fish begin to switch from feeding on invertebrates to eating fish (striped bass, shad, salmon, flatfish, herring, perch, bait). This trend continues in the subadults. Although adults are primarily piscivorous, striped bass continue to eat some invertebrates, such as crabs. The adults, and perhaps the subadults, reduce their feeding in the spring and early summer during the spawning season.

The life cycle of *Anisakis* sp. usually includes a marine crustacean as the first intermediate host, a fish or squid as the second host, and a marine mammal as the definitive host. The larva of *Anisakis* sp. is well known as a potential hazard to human health if

consumed in raw or improperly prepared seafood (Moser, Love, and Sakanari 1984). San Francisco Bay striped bass infected with larval *Anisakis* do not appear to be a major health hazard. Rarely were these larvae found in the muscle tissue. In addition, most of the larvae degenerate in the body cavity after a few weeks. The results of this study suggest that some degenerating larvae and the fibrous "rafts" can persist in striped bass for at least 33 wk. This slow turnover rate of larvae probably contributes to the increase in intensity of infection with time.

Wootten and Smith (1975) showed that trout can be serially infected with *Anisakis* sp. larvae from herring. Many of the fish consumed by striped bass are infected by *Anisakis* sp. larvae (Love and Moser 1983). The serial infection experiments with *Anisakis* sp. larvae may explain the increase in prevalence and intensity as striped bass become more piscivorous at 4 yr. Although numerous fibrous "rafts" were found in the mesenteries and next to the peritoneum of field fish, they did not form open lesions as described for the metacestodes of *L. tenuis* in San Francisco Bay striped bass (Moser, Sakanari, Wellings, and Lindstrom 1984).

The life cycle of *Lacistorhynchus tenuis*, although not completely described, most probably includes the adult in an elasmobranch, such as the leopard shark, *Triakis semifasciata*, common to San Francisco Bay, the procercoid in a crustacean, and the metacestode in fish. *Lacistorhynchus tenuis* metacestodes have been reported worldwide from more than 60 species of hosts (Moser 1983), including striped bass from the east coast and other San Francisco Bay fish, with no apparent harm to the host. However, Moser, Sakanari, Wellings, and Lindstrom (1984) have shown that these metacestodes are pathological to striped bass found in the San Francisco Bay-Delta area.

The sharp rise in prevalence of *L. tenuis* metacestodes in the 1-yr-olds may be related to the movement of this age class from the Delta area into San Francisco Bay where they are more likely to encounter the infected intermediate hosts. The continued increase in the intensity of metacestodes through time may reflect both the acquisition of new metacestodes and their persistence in the host. Preliminary studies (Sakanari, Moser) suggest that metacestodes of *L. tenuis* can be transfered serially from white croaker, *Genyonemus lineatus*, to striped bass. These studies suggest that the prevalence of metacestodes from the YOY to the 1 yr class may also be related to increased piscivory. The increased clumped distribution of parasites with the age of the host may indicate striped bass vary in their susceptibility to these parasites. Some hosts may eliminate the parasites before they can become embedded in the mesentaries or muscle tissue. Striped bass were introduced into San Francisco Bay from the east coast approximately 100 yr ago. The type of incompatibility shown here has not been described between east coast striped bass and their helminth parasites. Current studies are investigating the possibility that the west coast striped bass are immunologically incompatible with the indigenous strain of parasite.

ACKNOWLEDGMENTS

We thank the staffs at Marine Studies, University of California, Santa Cruz, and the National Marine Fisheries Service, Tiburon, CA; Marvin Jung, State Water Resources Control Board; Frank Fisher and the California Department of Fish and Game, Kinnetics Laboratory; Elmer Noble, University of California, Santa Barbara; James Berry, Scripps Institute of Oceanography, La Jolla, CA; Donald Gruber, Pinole, CA; Thomas Tucker, Steinhart Aquarium, California Academy of Sciences.

LITERATURE CITED

CAILLIET, G. M.
1976. Several approaches to the feeding ecology of fishes. First Pac. N.W. Tech. Workshop. Sess. 1. Workshop Proc., Astoria, Oreg., October 13-15, 1976.

EDWARDS, S. R., and F. M. NAHHAS.
1968. Some endoparasites of fishes from the Sacramento-San Joaquin Delta, California. Calif. Fish Game 54:247-256.

HENSLEY, G. H., and F. M. NAHHAS.
1975. Parasites of fishes from the Sacramento-San Joaquin Delta, California. Calif. Fish Game 61:201-208.

LOVE, M. S., and M. MOSER.
1983. A checklist of parasites of California, Oregon, and Washington marine and estuarine fishes. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-777, 576 p.

MORIGUCHI, M. N.
1973. Food habits of the striped bass, *Roccus saxatilis*, (Walbaum) in San Francisco Bay with notes on the major areas of the fishery. M.A. Thesis, San Francisco State Univ., San Francisco, CA, 61 p.

MOSER, M.
1983. Parasites of San Francisco Bay-Delta area stripped bass (*Morone saxatilis*) and other San Francisco Bay fishes. Final Rep., Calif. State Water Res. Control Board, Sacramento, CA, 114 p.

MOSER, M., M. LOVE, and J. SAKANARI.
1984. Common parasites of California fishes. Calif. Dep. Fish Game, 20 p.

MOSER, M., J. SAKANARI, S. WELLINGS, and K. LINDSTROM.
1984. Incompatibility between San Francisco striped bass, *Morone saxatilis* (Walbaum), and the metacestode, *Lacistorhynchus tenuis* (Beneden 1858). J. Fish Dis. 7:397-400.

MOYLE, P. B.
1976. Inland fishes of California. Univ. Calif. Press, Berkeley, 405 p.

ORSI, J. J.
1971. The 1965-1967 migrations of the Sacramento-San Joaquin Estuary striped bass population. Calif. Fish Game 57:257-267.

ORSI, J. J., and A. C. KNUTSON.
1979. The role of mysid shrimp in the Sacramento-San Joaquin estuary and factors affecting their abundance and distribution. *In* T. J. Conomos (editor), San Francisco Bay: The urbanized estuary, p. 401-408. Pac. Div./A.A.A.S., San Francisco State Univ., CA, June 12-16, 1977.

SOKAL, R. R., and F. J. ROHLF.
1969. Biometry. The principles and practice of statistics in biological research. W. H. Freeman and Co., San Francisco, CA, 776 p.

TANAKA, J., and M. TORISU.
1978. *Anisakis* and eosinophil. I. Detection of a soluble factor selectively chemotactic for eosinophils in the extract from *Anisakis* larvae. J. Immunol. 120:745-749.

THOMAS, J. L.
1967. The diet of juvenile and adult striped bass, *Roccus saxatilis*, in the Sacramento-San Joaquin River system. Calif. Fish Game 53:49-62.

TURNER, J. L., and D. W. KELLEY.
1966. Ecological studies of the Sacramento-Joaquin Delta. Calif. Dep. Fish Game, Fish Bull. 136, Part II, Fishes of the Delta, 168 p.

WOOTTEN, R., and J. W. SMITH.
1975. Observational and experimental studies on the acquisition of *Anisakis* sp. larvae (Nematoda: Ascaridida) by trout in fresh water. Int. J. Parasitol. 5:373-378.

NOAA TECHNICAL REPORT NMFS

Guidelines for Contributors

CONTENTS OF MANUSCRIPT

First page. Give the title (as concise as possible) of the paper and the author's name, and footnote the author's affiliation, mailing address, and ZIP code.

Contents. Contains the text headings and abbreviated figure legends and table headings. Dots should follow each entry and page numbers should be omitted.

Abstract. Not to exceed one double-spaced page. Footnotes and literature citations do not belong in the abstract.

Text. See also Form of the Manuscript below. Follow the *U.S. Government Printing Office Style Manual,* 1973 edition. Fish names, follow the American Fisheries Society Special Publication No. 12, *A List of Common and Scientific Names of Fishes from the United States and Canada,* fourth edition, 1980. Use short, brief, informative headings in place of "Materials and Methods."

Text footnotes. Type on a separate sheet from the text. For unpublished or some processed material, give author, year, title of manuscript, number of pages, and where it is filed—agency and its location.

Personal communications. Cite name in text and footnote. Cite in footnote: John J. Jones, Fishery Biologist, Scripps Institution of Oceanography, La Jolla, CA 92037, pers. commun. 21 May 1977.

Figures. Should be self-explanatory, not requiring reference to the text. All figures should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Photographs and line drawings should be of "professional" quality—clear and balanced, and can be reduced to 42 pices for page width or to 20 picas for a single-column width, but no more than 57 picas high. Photographs and line drawings should be printed on glossy paper—sharply focused, good contrast. Label each figure. DO NOT SEND original figures to the Scientific Editor; NMFS Scientific Publications Office will request these if they are needed.

Tables. Each table should start on a separate page and should be self-explanatory, not requiring reference to the text. Headings should be short but amply descriptive. Use only horizontal rules. Number table footnotes consecutively across the page from left to right in Arabic numerals; and to avoid confusion with powers, place them to the *left* of the numerals. If the original tables are typed in our format and are clean and legible, these tables will be reproduced as they are. In the text all tables should be cited consecutively and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page.

Acknowledgments. Place at the end of text. Give credit only to those who gave exceptional contributions and *not* to those whose contributions are part of their normal duties.

Literature cited. In text as: Smith and Jones (1977) or (Smith and Jones 1977); if more than one author, list according to years (e.g., Smith 1936; Jones et al 1975; Doe 1977). All papers referred to in the text should be listed alphabetically by the senior author's surname under the heading "Literature Cited"; only the author's surname and initials are required in the author line. The author is responsible for the accuracy of the literature citations. Abbreviations of names of periodicals and serials should conform to *Biological Abstracts List of Serials with Title Abbreviations.* Format, see a recent Technical Report.

Abbreviations and symbols. Common ones, such as mm, m, g, ml, mg, °C (for Celsius), %, ‰, etc., should be used. Abbreviate units of measures only when used with numerals; periods are rarely used in these abbreviations. But periods are used in et al., vs., e.g., i.e., Wash. (WA is used only with ZIP code), etc. Abbreviations are acceptable in tables and figures where there is lack of space.

Measurements. Should be given in metric units. Other equivalent units may be given in parentheses.

FORM OF THE MANUSCRIPT

Original of the manuscript should be typed double-spaced on white bond paper. Triple space above headings. Send good duplicated copies of manuscript rather than carbon copies. The sequence of the material should be:

FIRST PAGE
CONTENTS
ABSTRACT
TEXT
LITERATURE CITED
TEXT FOOTNOTES
APPENDIX
TABLES (provide headings, including "Table" and Arabic numeral, e.g., Table 1.—, Table 2.—, etc.)
LIST OF FIGURE LEGENDS (entire legend, including "Figure" and Arabic numeral, e.g., Figure 1.—, Figure 2.—, etc.)
FIGURES

ADDITIONAL INFORMATION

Send ribbon copy and two duplicated copies of the manuscript to:

Dr. William J. Richards, Scientific Editor
Southeast Fisheries Center Miami Laboratory
National Marine Fisheries Service, NOAA
75 Virginia Beach Drive
Miami, FL 33149

Copies. Fifty copies will be supplied to the senior author and 100 to his organization free of charge.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS OFFICE
7600 SAND POINT WAY N.E.
BIN C15700
SEATTLE, WA 98115

OFFICIAL BUSINESS

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161